

The Burgers equation as electrohydrodynamic model in plasma physics

E. Moreau and O. Vallée

Laboratoire d'Analyse Spectroscopique et d'Énergétique des Plasmas

Faculté des Sciences, rue Gaston Berger BP 4043

18028 Bourges Cedex France

(Dated: February 2, 2008)

Abstract

It is shown that the one dimensional Poisson equation governing the electric field in a plasma, yields to an inhomogeneous Burgers type equation which may be solved analytically for some particular cases. The solution obtained for these cases shows an elastic behaviour of the associated electric force in the long time limit.

PACS numbers: 47.65.+a, 51.50.+v, 52.30.-q

This paper points out the possibility to treat analytically some problems occurring in electro or magnetohydrodynamics (EMHD) models. It is well known that the equations describing fluid models are difficult to handle because of their mathematical complexities: most of them are partial differential equations and present nonlinear properties [1], thereby, the analytic approach is nearly totally neglected. Nevertheless, there exist some situations where analytic solutions may be obtained. One famous example is the Burgers equation [2, 3], and especially the homogeneous Burgers equation, which is known to give valuable results in turbulence simulations. Moreover, this approximation of the Navier-Stokes equation presents some cases which are integrable. In particular, the homogeneous Burgers equation in the inviscid limit has been studied a lot over past decades, and has given rise to the “Burgulence approach” [4]. But, for a real EMHD problem, electric and magnetic fields appear by the way of Maxwell equations, so, the problem presents additional difficulties due to the non linear coupling between fluid equations and Maxwell equations. A first approach in the resolution of such equations has been proposed by some authors like Thomas [5] and more recently by Olesen [6]. In these works, a 1+1 dimensional MHD problem is treated with particular configuration for the velocity and the magnetic field. The motion equation

$$\partial_t \mathbf{v} + (\mathbf{v} \nabla) \mathbf{v} = \nu \nabla^2 \mathbf{v} + (\nabla \times \mathbf{B}) \times \mathbf{B}, \quad (1)$$

is thus coupled with an equation of evolution for the magnetic field

$$\partial_t \mathbf{B} = \nabla \times (\mathbf{v} \times \mathbf{B}) + (1/\sigma) \nabla^2 \mathbf{B}, \quad (2)$$

to give a homogeneous Burgers type equation which may consequently be integrated. In this model, only the magnetic field is considered, but this allows to take into account the Lorentz force acting on a velocity field. So, these works are based on a nonlinear connection between an equation of motion where appears a magnetic field (Eq. (1)), and a specific equation for the magnetic field (Eq. (2)). But, another interesting aspect of a EMHD problem, consists to study the influence of an electric field on a velocity field. Thus, the present paper may be considered as a complementary approach facing Olesen’s works, in the sense where we propose an electric version of Eqs. (1) and (2). In ref.[7] we have solved a Burgers equation with an elastic forcing term, which allows to treat a velocity field undergoing a fluctuating strength force. Hence, the problem will consist to show that an electric field may be considered as an elastic term of force for some physical problems. This will be done in the following by using the Poisson equation. In order to reach this result, we propose an analytic resolution of a one dimensional problem encountered in ionised media, when

the electric field is dominant in front of the magnetic field. We may have this situation for example in electric discharges or electric arcs studies. For this purpose, we consider a constant electric field E_a applied to a plasma in which we assume the existence of a constant electron flow (created for example by electrodes system and depending of E_a), which may be interpreted as a source term. Moreover, the plasma is assumed to be composed of electrons and motionless ions, in such a way that we can write the one dimensional Poisson equation as

$$\partial_x E = -\frac{|e|}{\epsilon_0}n, \quad (3)$$

where n is the electronic density, and $|e|$ the absolute value of the electron charge. If the diffusion process is taken into account, then the electronic current density will read

$$J = -\mu n(E - E_a) - \nu \partial_x n; \quad (4)$$

with μ standing for the (constant) electronic mobility, and ν the electronic diffusion coefficient. After what, we use the equation of conservation for the electronic density:

$$\partial_t n + \partial_x J = S, \quad (5)$$

where S represents the constant source term. From this, combining relations (3), (4) and (5) in order to remove the density, we have successively

$$J = \frac{\mu\epsilon_0}{|e|}(E - E_a)\partial_x E + \frac{\nu\epsilon_0}{|e|}\partial_{xx}E, \quad (6)$$

and

$$-\frac{\epsilon_0}{|e|}\partial_x \partial_t E + \partial_x \left(\frac{\mu\epsilon_0}{|e|}(E - E_a)\partial_x E + \frac{\nu\epsilon_0}{|e|}\partial_{xx}E \right) = S; \quad (7)$$

which may be integrated to give

$$\partial_t E - \mu(E - E_a)\partial_x E = \nu\partial_{xx}E - \frac{|e|}{\epsilon_0}xS + c(t), \quad (8)$$

$c(t)$ being an arbitrary function. After what, carrying out the following change of variables

$$x = \mu\bar{x}, \quad \bar{\nu} = \frac{\nu}{\mu^2}, \quad \bar{S} = S\frac{|e|\mu}{\epsilon_0};$$

Eq. (8) reads

$$\partial_t E - (E - E_a)\partial_{\bar{x}}E = \bar{\nu}\partial_{\bar{x}\bar{x}}E - \bar{S}\bar{x} + c(t). \quad (9)$$

Thus, the quantity E_a being constant, a Burgers type equation may be obtained for the variable $E - E_a$:

$$\partial_t(E - E_a) - (E - E_a)\partial_{\bar{x}}(E - E_a) = \bar{\nu}\partial_{\bar{x}\bar{x}}(E - E_a) - \bar{S}\bar{x} + c(t). \quad (10)$$

The local field $E(\bar{x}, t)$ may then be determined by putting

$$\mathcal{E}(\bar{x}, t) = E(\bar{x}, t) - E_a, \quad (11)$$

that is to say that we have to solve the following one dimensional Burgers equation with an elastic forcing term

$$\partial_t \mathcal{E} - \mathcal{E}\partial_{\bar{x}} \mathcal{E} = \bar{\nu}\partial_{\bar{x}\bar{x}} \mathcal{E} - \bar{x}\bar{S} + c(t); \quad (12)$$

with

$$\mathcal{E}(\bar{x}, 0) = -E_a \quad (13)$$

as initial condition if initially there is no charges separation. Notice that a minus sign appears with the non linear term, with the meaning that the electrons spread in the opposite direction of the electric field. Then, We have shown ref.[7] the possibility to solve analytically such an equation by two equivalent methods. Putting $c(t)$ to zero, it is hence possible to show that the solution reads, in physical coordinates (x, t) ,

$$\mathcal{E}(x, t) = -\frac{\gamma}{\mu} (1 - e^{-2\gamma t}) x - E_a e^{-\gamma t} \quad (14)$$

where we have put

$$\gamma = \sqrt{\frac{S|e|\mu}{\epsilon_0}}. \quad (15)$$

It must be underlined that the effective electric field acting on any electron of the system is in fact $-\mathcal{E}(x, t)$. With this in mind, a physical interpretation of the problem may be obtained in the asymptotic mode when $S \geq 0$ (i.e $\gamma \in \mathbb{R}^+$). In this case, the asymptotic limit, reached for $t \gg 1/\gamma$, simply reads

$$\lim_{t \rightarrow \infty} \mathcal{E}(x, t) = -\frac{\gamma}{\mu} x, \quad (16)$$

which is a stationary solution. In other words, the system has reached a stationary state. Thus, we can deduce that the electric force F acting on an electron is related to an elastic forcing term. As a matter of fact, we have in this stationary state

$$F = -|e| \times -\lim_{t \rightarrow \infty} \mathcal{E}(x, t) = -|e|\frac{\gamma}{\mu} x = -\kappa^2 x, \quad (17)$$

where κ^2 stands for a string constant and is given by

$$\kappa^2 = |e| \frac{\gamma}{\mu} = \left(\frac{S|e|^3}{\mu\epsilon_0} \right)^{1/2}. \quad (18)$$

Notice that an electron undergoing the single force (17) (low viscosity), will oscillate with a pulsation ω such as

$$\omega^2 = \frac{\kappa^2}{m_e} = \frac{|e|\gamma}{\mu m_e}; \quad (19)$$

m_e standing for the electronic mass. This expression has to be compared with the plasma frequency ω_p :

$$\omega_p^2 = \frac{|e|^2 n}{\epsilon_0 m_e}, \quad (20)$$

with n given by the Poisson equation (3). In the long time limit we have

$$n = -\frac{\epsilon_0}{|e|} \partial_x \lim_{t \rightarrow \infty} E(x, t) = \frac{\epsilon_0 \gamma}{|e| \mu}. \quad (21)$$

So,

$$\omega_p^2 = \frac{|e|\gamma}{\mu m_e} = \omega^2. \quad (22)$$

Hence, for a low viscosity system, the electrons of our model oscillate with the theoretical plasma frequency. In a more general case, when viscosity effects have to be taken into consideration, we assume the following motion equation satisfied for the electronic velocity u

$$\partial_t u + u \partial_x u = \nu \partial_{xx} u - \kappa^2 x. \quad (23)$$

In this case, an electron undergoes both viscosity and electric field. The Eq. (23) is an inhomogeneous Burgers equation with an elastic term of force. But, we have shown ref.[7] that in such a case, a physical solution was

$$\lim_{t \rightarrow \infty} u(x, t) = \kappa x. \quad (24)$$

Then, assuming that the velocity u writes (relation (4))

$$u(x, t) = -\mu \mathcal{E}(x, t) - \nu \partial_x \ln[n(x, t)], \quad (25)$$

with $n(x, t)$ given by the Poisson equation (3), we have in the asymptotic limit

$$\lim_{t \rightarrow \infty} u(x, t) = -\mu \lim_{t \rightarrow \infty} \mathcal{E}(x, t) - \nu \lim_{t \rightarrow \infty} \partial_x \ln[n(x, t)]. \quad (26)$$

Then, a relation connecting the source term with the electronic mobility appears from the limits (16) and (24):

$$S\mu^3 = |e|\epsilon_0 \iff \mu \propto S^{-1/3}. \quad (27)$$

This result may have interesting consequences when the source term is known (for example the number of electrons created by an electrode per unit of time determined by the Richardson-Dushman law), because it permits to determine the electronic mobility. Conversely, since the mobility is considered as a gas property, the relation (27) gives a first approximation of the value of the source term.

In summary, we have presented an electrohydrodynamical model describing the evolution of a local electric field in a plasma. Since we make the assumption of motionless ions in the model, we have shown the possibility to derive a new equation of evolution for the electric field obeying to the Poisson equation. Thereby, the results presented here have to be seen as a continuation of Olesen's works, in the sense where we have presented and solved an electric version of Eqs. (1) and (2). The two basic equations being the equation of motion

$$\partial_t u + u \partial_x u = \nu \partial_{xx} u + |e| \mathcal{E}, \quad (28)$$

and the Poisson equation

$$\partial_x \mathcal{E} = -\frac{|e|}{\epsilon_0} n. \quad (29)$$

Then, we have found a solution for the electric field of the form

$$\mathcal{E}(x, t) = E(x, t) - E_a = A(t)x + B(t), \quad (30)$$

with the properties

$$\lim_{t \rightarrow \infty} A(t) = -\frac{\gamma}{\mu} \quad \text{and} \quad \lim_{t \rightarrow \infty} B(t) = 0. \quad (31)$$

with the meaning that the electric field behaves like an oscillating force in the long time limit, and with a string constant proportional to the square root of the source term. Furthermore, a relation may be obtained, still in the asymptotic mode, between the term of source S and the electronic mobility μ , and may be interpreted as a general scale law under some hypothesis. Notice that we have focused our attention on a positive term of source. The case of a negative S , would give rise to another kind of physics which may be interesting to treat in a forthcoming paper.

As a corollary we enforce the result discussed in ref.[7], which states that the transformation

$$u(x, t) = -2\nu \frac{1}{P(x, t)} \partial_x P(x, t) - \kappa x, \quad (32)$$

introduced in the Burgers equation (23), yields to the Fokker-Planck equation

$$\partial_t P(x, t) = \kappa \partial_x (x P(x, t)) + \nu \partial_{xx} P(x, t). \quad (33)$$

And this expression may be considered as a collisional term in a Vlasov-Poisson model (without magnetic field) ; the electric field being determined by the method presented here. Under specific assumptions, this remark would open an interesting way of investigation.

- [1] M. LESIEUR, *turbulence in Fluids*, Fluid Mechanics and its Applications, Vol. 40, Kluwer, 1997.
- [2] J. M. BURGERS, Adv. Appl. Mech. **1** (1949) 171.
- [3] J. M. BURGERS, *The non linear diffusion equation* (Reidel, Boston, 1974).
- [4] U. FRISCH, & J. BEC, LES HOUCHEs 2000: *New Trends in Turbulence* M. Lesieur, ed., Springer EDP-Sciences (2001), nlin.cd/0012033.
- [5] J. H. THOMAS, Phys. Fluids, **11**, 1245 (1968).
- [6] P. OLESEN, ARXIV:NLIN.SI/ 0304005 version 2.
- [7] E. MOREAU & O. VALLÉE. *Connection between the Burgers equation with an elastic forcing term and a stochastic process*; arXiv:physics/0412136 .